a hydroxyl group, the molar ratio of R<sub>3</sub><sup>1</sup>1SiO<sub>4</sub> unit to SiO2 unit being in the range of from 0.5 to 1.2.

- (C) an organohydrogenpolysiloxane containing at least two hydrogen atoms attached to silicon atoms in a molecule, in an amount such that the ratio of the total molar number of hydrogen atoms in component (C) and the total molar number of vinvl groups in components (A) and (E) is 0.1 to 5,
- (D) a catalytic amount of a platinum catalyst, and (E) an organopolysiloxane of the following general formula (3);

$$(CH_2=CH_{\frac{1}{p_n}}^{R_{3-n}} \underset{Si \leftarrow OSi}{\overset{R}{\underset{p}{\downarrow}}} \underset{Q}{\overset{R}{\underset{q}{\downarrow}}} \underset{OSi \leftarrow CH}{\overset{R_{3-n}}{\underset{q}{\downarrow}}} \underset{OSi \leftarrow CH=CH_2}{\overset{(3)}{\underset{n}{\downarrow}}} 1$$

wherein R and n are as defined above, p is an integer of from 0 to 1,000, q is an integer of from 10 to 20 1,000, and  $0 \le p/q \le 1$ .

- 2. The composition of claim 1 which contains about 100 to about 200 parts of component (B).
- 3. The composition of claim 1 wherein component 25 (C) is added such that the ratio of the total molar number of hydrogen atoms in component (C) and the total molar number of vinyl groups in components (A) and (E) ranges from 0.1 to 5.
- 4. The composition of claim 1 which contains 100 30 parts of component (A) and about 0.2 to about 10 parts of component (E).
- 5. The composition of claim 1, wherein 1-5 parts by weight of (E) per 100 parts by weight of (A) are used.
- 6. The composition of claim 1, wherein 1-4 parts by weight of (E) per 100 parts by weight of (A) are used.
- 7. A silicone self-adhesive composition comprising 100 parts by weight of at least one organopolysiloxane anes of the following general formulae (1) and (2);

## -continued

wherein R is a monovalent hydrocarbon group excluding alkenyl, letter n is an integer of from 0 to 3, k and m 10 each are 0 or a positive integer, with the proviso that m is an integer of at least 2 when n is 0, and j and 1 each are an integer of at least 2,000,

- (B) an organopolysiloxane comprising R<sub>3</sub><sup>1</sup>SiO<sub>4</sub> and SiO<sub>2</sub> units wherein R<sup>1</sup> is a monovalenthydrocarbon group or a hydroxyl group, the molar ratio of R<sub>3</sub><sup>1</sup>1SiO<sub>4</sub> unit to SiO<sub>2</sub> unit being in the range of from 0.5 to 1.2,
- (C) an organohydrogenpolysiloxane containing at least two hydrogen atoms attached to silicon atoms in a molecule, in an amount such that the ratio of the total molar number of hydrogen atoms in component (C) and the total molar number of vinyl groups in components (A) and (E) is 0.1 to 5,
- (D) a catalytic amount of a platinum catalyst, and (E) 1-5 parts by weight of an organopolysiloxane of the following general formula (3);

$$(CH2=CH)_{\overline{n}}^{R_{3-n}} \begin{array}{cccc} R & R & R_{3-n} & (3) \\ \vdots & \vdots & \vdots & \vdots \\ R & CH=CH2 & OSi+CH=CH2)_{n} \\ R & CH=CH2 & CH=CH2 & (3) \\ \vdots & \vdots & \vdots & \vdots \\ R & CH=CH2 & (3) \\ \vdots & \vdots & \vdots \\ R &$$

wherein R and n are as defined above, p is an integer of from 0 to 1,000, q is an integer of from 10 to 1,000, and  $0 \le p/q \le 1$ .

- 8. The composition of claim 7, wherein 100-250 parts by weight of (B) per 100 parts by weight of (A) are used.
- 9. The composition of claim 7, wherein 100-200 parts selected from the group consisting of organopolysilox- 40 by weight of (B) per 100 parts by weight of (A) are
  - 10. The composition of claim 7, wherein 1-4 parts by weight of (E) per 100 parts by weight of (A) are used.
  - 11. The composition of claim 7, wherein 1-3 parts by 45 weight of (E) per 100 parts by weight of (A) are used.

## 60